

Supporting Information for

Na₂Fe(C₂O₄)F₂: A new iron-based polyoxyanion cathode for Li/Na ion batteries

Wenjiao Yao,[†] Moulay-Tahar Sougrati,^{§,‡} Khang Hoang,[¶] Jianing Hui,[†] Philip Lightfoot,^{†,*} Anthony Robert Armstrong^{†,‡*}

[†] School of Chemistry, University of St Andrews, St Andrews, Fife KY16 9ST, UK

[§] Université de Montpellier 2 Place Eugène Bataillon - CC 1502, 34095 Montpellier CEDEX 5, France

[¶] Center for Computationally Assisted Science and Technology, North Dakota State University, Fargo, ND 58108, USA

[‡]ALISTORE-ERI, 80039, Amiens Cedex, France

*P. Lightfoot: pl@st-andrews.ac.uk

*A. R. Armstrong: ara@st-andrews.ac.uk

Table of Contents

1. Unit cell information, atomic coordinates and Bond valence sums (BVS) for metal atoms in $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$. Table S1
2. Selected bond lengths in $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$. Table S2
3. Coordination polyhedra of $[\text{FeO}_4\text{F}_2]$, $[\text{NaO}_x\text{F}_y]$ in $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$. Figure S1
4. Morphology characterization of hand ground $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$ powder sample. Figure S2
5. UV-Vis Absorption Spectrum for $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$ powder. Figure S3
6. dQ/dV curve of initial charge/discharge profiles in LIB. Figure S4
7. Mössbauer parameters for $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$ at different states corresponding to Figure 4d-f. Table S3
8. Assignment of IR spectrum peaks.
9. Solubility analysis. PXRD of left-over sample from $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$ immersed in water for five days. Figure S5
10. Analysis of left-over sample from $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$ immersed in 1M LiBr in CH_3CN . Figure S6

1 Table S1 Unit cell information and atomic coordinates and BVS for metal atoms in Na₂Fe(C₂O₄)F₂

Atom	Wyck.	Site	x/a	y/b	z/c	Uequiv[Å ²]	BVS
Fe1	8f	1	0.17403(2)	0.16083(5)	0.21291(2)	0.00717(14)	2.022
Na1	8f	1	0.29790(6)	0.00423(16)	0.51447(8)	0.0121(2)	1.104
Na2	8f	1	0.06659(6)	-0.03217(16)	0.40099(9)	0.0116(2)	1.022
F1	8f	1	0.18763(9)	0.1500(2)	0.06049(11)	0.01252(3)	
F2	8f	1	0.17823(9)	0.1726(2)	0.37587(11)	0.01175(3)	
O1	8f	1	-0.05003(10)	0.5774(3)	0.10778(14)	0.00857(3)	
O2	8f	1	0.31304(10)	0.0289(3)	0.28934(13)	0.00644(3)	
O3	8f	1	0.39911(12)	0.3347(3)	0.31129(17)	0.01257(4)	
O4	8f	1	0.0364(1)	0.2707(3)	0.13864(14)	0.00855(3)	
C1	8f	1	0.38838(15)	0.1302(4)	0.31611(19)	0.00981(4)	
C2	8f	1	0.02406(15)	0.4791(4)	0.13964(18)	0.01041(4)	

2 Table S2 Selected bond lengths in Na₂Fe(C₂O₄)F₂

Atom1	Atom 2	Distance (Å)	Atom 1	Atom 2	Distance (Å)
Fe1	F1	1.9633(15)	Na2	F2	2.2624(18)
	F2	1.9782(15)		F1	2.2791(14)
	O4	2.1268(15)		O1	2.340(2)
	O2	2.1907(15)		O4	2.3587(19)
	O2	2.2051(18)		O3	2.586(2)
	O3	2.2255(18)	C1	O3	1.235(3)
Na1	F1	2.2253(14)		O2	1.266(3)
	F2	2.2543(19)		C2	1.570(3)
	O1	2.2987(17)	C2	O1	1.237(3)
	F2	2.2994(16)		O4	1.258(3)
	F1	2.3490(16)		C1	1.570(3)

3 Coordination polyhedra in $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$




Figure S1 (a) FeO_4F_2 (b) NaOF_4 (c) NaO_3F_2

4 Morphology characterization of hand ground $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$ powder sample




Figure S2 SEM image of hand ground $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$ powder

5 UV-Vis Absorption Spectrum for $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$ powder




Figure S3 UV-Vis Absorption spectrum for $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$ powder.

6 dQ/dV curve of initial charge/discharge profiles in LIB




Figure S4 dQ/dV curve of initial charge/discharge profile in LIB

7 Table S3 Mössbauer parameters for Na₂Fe(C₂O₄)F₂ at different states corresponding to Figure 4d-f.

<i>spectrum</i>	<i>Comp.</i>	<i>IS mm/s</i>	<i>QS mm/s</i>	<i>LW mm/s</i>	<i>Abs %</i>	<i>Attribution</i>
Pure (Figure 4d)	Green-Na ₂ Fe(C ₂ O ₄)F ₂	1.23	2.56	0.28	91	Fe ²⁺
	Blue- [†]	1.22	1.58	0.63	9	Fe ²⁺
4.3 V- Charged (Figure 4e)	Green- Na ₂ Fe(C ₂ O ₄)F ₂	1.08	2.16	0.31	2	Fe ²⁺
	Blue-	1.24	2.76	0.27	8	Fe ²⁺
	Red-NaFe(C ₂ O ₄)F ₂	0.41	0.55	0.55	82	Fe ³⁺
	Pink-NaFe(C ₂ O ₄)F ₂	0.43	1.62	0.27	8	Fe ³⁺
2.0 V- Discharged (Figure 4f)	Green-Na ₂ Fe(C ₂ O ₄)F ₂	1.23	2.57	0.26	69	Fe ²⁺
	Blue-	1.20	1.93	0.45	5	Fe ²⁺
	Red-NaFe(C ₂ O ₄)F ₂	0.42	0.57	0.64	26	Fe ³⁺

[†]The blue line in the pristine sample indicates a minor environment, also in Fe(II) (high spin) but with slightly different parameters. It suggests a contribution arising from some defects in the bulk material, very small particles and/or amorphous material, or could be an impurity, such as amorphous Fe(C₂O₄)• 2H₂O.

8 IR Spectroscopy of $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$.

The absorption peaks in the IR spectrum are assigned as:

- 434 cm^{-1} : Fe-F antisymmetric stretching;
- 503 cm^{-1} : Fe-O symmetric stretching in the $[\text{FeO}_4]$ plane;
- 783 cm^{-1} : $\delta(\text{OCO})$ bending modes;
- 902 cm^{-1} : $\nu(\text{C-C})$ stretching vibration;
- 1309 cm^{-1} : $\nu_s(\text{COO})$ involving dangling oxygen;
- 1456 cm^{-1} : $\nu_s(\text{COO})$;
- 1606 cm^{-1} : $\nu_{as}(\text{COO})$ involving dangling oxygen;
- 1674 cm^{-1} : $\nu_{as}(\text{COO})$.

9 Solubility Analysis




Figure S5. PXRD of left-over sample from $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$ immersed in water for five days.
Red: $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$, Blue: $\text{Fe}(\text{C}_2\text{O}_4) \cdot 2\text{H}_2\text{O}$

10 Analysis of left-over sample from $\text{Na}_2\text{Fe}(\text{C}_2\text{O}_4)\text{F}_2$ immersed in 1M LiBr in CH_3CN

(a) EDX analysis

Spectrum processing:

Peak possibly omitted: 1.500 keV

Processing option: All elements analyzed (Normalised)

Number of iterations = 7


Standard:

C CaCO_3 1-Jun-1999 12:00 AM

O SiO_2 1-Jun-1999 12:00 AM

F MgF_2 1-Jun-1999 12:00 AM

Fe Fe 1-Jun-1999 12:00 AM



Element	Weight%	Atomic%
C K	6.98	10.94
O K	11.84	13.94
F K	72.96	72.35
Fe K	8.23	2.78
Totals	100.00	




Figure S6. (a) SEM image and EDX spectrum

(b) PXRD analysis




Figure S6. (b) PXRD of left-over sample